

Detailed Studies of Cloud Cores: Probing the Initial Conditions for Protostellar Collapse

Ph. André¹, A. Belloche², P. Hennebelle³ and D. Ward-Thompson⁴

¹ *CEA Saclay, Service d’Astrophysique, Orme des Merisiers, Bât. 709, F-91191 Gif-sur-Yvette Cedex, France*

² *Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*

³ *Laboratoire de Radioastronomie, ENS, 24 rue Lhomond, F-75231 Paris Cedex 05, France*

⁴ *Dept of Physics, Cardiff University, PO Box 913, Cardiff, UK*

Received December 4, 2003

Abstract. Improving our understanding of the earliest stages of star formation is crucial to gain insight into the origin of stellar masses, multiple systems, and protoplanetary disks. We discuss recent advances made in this area thanks to detailed mapping observations at infrared and (sub)millimeter wavelengths. Although ambipolar diffusion appears to be too slow to play a direct role in the formation of dense cores, there is nevertheless good evidence that the gravitational collapse of isolated protostellar cores is strongly magnetically controlled. We also argue that the beginning of protostellar collapse is much more violent in cluster-forming clouds than in regions of distributed star formation.

Key words: ISM: structure – ISM: gravitational collapse – Stars: protostellar accretion rates

1. INTRODUCTION

One of the main limitations in our present understanding of the star formation process is that we do not know well the initial conditions for protostellar collapse. In particular, there is a major controversy at the moment between two schools of thought for the formation and evolution of dense cores within molecular clouds: The classical picture based on magnetic support and ambipolar diffusion (e.g. Shu et al. 1987, 2003; Mouschovias & Ciolek 1999) has been seriously challenged by a new, more dynamical picture,

which emphasizes the role of supersonic turbulence in supporting clouds on large scales and generating density fluctuations on small scales (e.g. Klessen et al. 2000; Padoan & Nordlund 2002). Improving our knowledge of the earliest stages of star formation is thus of prime importance, especially since there is now good evidence that these stages control the origin of the stellar initial mass function (e.g. Motte, André, & Neri 1998; Testi & Sargent 1998; Johnstone et al. 2000).

Here, we discuss recent observational advances concerning the density and velocity structure of cloud cores (§2 and §3), and we compare the results with the predictions of theoretical models (§4).

2. DENSITY STRUCTURE OF PRESTELLAR CORES

Two techniques have been used to trace the density structure of cloud cores (see Fig. 1a): (1) mapping the optically thin (sub)millimeter continuum *emission* from the cold dust contained in the cores, and (2) mapping the same cold core dust in *absorption* against the background infrared emission (originating from warm cloud dust or remote stars).

Ward-Thompson et al. (1994, 1999) and André et al. (1996) employed the first approach to probe the structure of prestellar cores (see also Shirley et al. 2000). Under the simplifying assumption of spatially uniform dust temperature and emissivity properties, they concluded that the radial density profiles of isolated prestellar cores were *not* consistent with the single $\rho(r) \propto r^{-2}$ power law of the singular isothermal sphere (SIS) but were flatter than $\rho(r) \propto r^{-1}$ in their inner regions (for $r \leq R_{flat}$), and approached $\rho(r) \propto r^{-2}$ only beyond a typical radius $R_{flat} \sim 2500\text{--}5000$ AU.

More recently, the use of the *absorption* approach, both in the mid-IR from space (e.g. Bacmann et al. 2000, Siebenmorgen & Krügel 2000) and in the near-IR from the ground (e.g. Alves et al. 2001), made it possible to confirm and extend the (sub)millimeter emission results, essentially independently of any assumption about the dust temperature distribution.

The typical column density profile found by these emission and absorption studies of prestellar cores has the following characteristics (see, e.g., Fig. 1b):

a) a flat inner region (of radius $R_{flat} = 5000 \pm 1000$ AU for L1689B according to the fitting analysis shown in Fig. 1b), b) a region roughly consistent with $N_{H_2} \propto \bar{r}^{-1}$ (corresponding to $\rho \propto r^{-2}$ for a spheroidal




Fig. 1. (a) ISOCAM $6.75\,\mu\text{m}$ absorption image of the prestellar core L1689B in Ophiuchus (scale on the right in MJy/sr). The $1.3\,\text{mm}$ continuum emission map of Andr et al. (1996) with the IRAM 30 m telescope is superposed as contours (levels: 10, 30, 50 mJy/ $13''$ -beam). (b) Column density profile of L1689B (crosses) derived from the absorption map shown in (a) by averaging the intensity over elliptical annuli for a 40° sector in the southern part of the core. The dashed curves show the most extreme profiles compatible with the data given the uncertainties affecting the absorption analysis. The solid curve is the best fit of a Bonnor-Ebert sphere model (embedded in a medium of uniform column density), obtained with the following parameters: $\rho_c/\rho_{out} = 40 \pm 15$ (i.e., well into the unstable regime), $T_{eff} = 50 \pm 20\,\text{K}$, $P_{ext}/k_B = 5 \pm 3 \times 10^5\,\text{K cm}^{-3}$. For comparison, the dotted line shows the $N_{\text{H}_2} \propto \bar{r}^{-1}$ profile of a SIS at $T = 10\,\text{K}$. (Adapted from Bacmann et al. 2000.)

core), c) a sharp edge where the column density falls off more rapidly than $N_{\text{H}_2} \propto \bar{r}^{-2}$ with projected radius defining the core outer radius ($R_{out} = 28000 \pm 1000\,\text{AU}$ for L1689B).

2.1. Comparison with models of core structure

The results summarized above set strong constraints on the density structure at the onset of protostellar collapse. The circularly-averaged column density profiles can often be fitted remarkably well with models of pressure-bounded Bonnor-Ebert spheres, as first demonstrated by Alves et al. (2001) for B68. This is also the case of L1689B as illustrated in Fig. 1b. The quality of such fits shows that Bonnor-Ebert spheroids provide a good, first order model for the structure of isolated prestellar cores. In detail, however, there are several prob-

lems with these Bonnor-Ebert models. First, the inferred density contrasts (from center to edge) are generally larger (i.e., $\gtrsim 20$ –80 – see Fig. 1b) than the maximum contrast of ~ 14 for stable Bonnor-Ebert spheres. Second, the effective core temperature needed in these models (for thermal pressure gradients to balance self-gravity) is often significantly larger than both the average dust temperature measured with ISOPHOT (e.g. Ward-Thompson et al. 2002) and the gas temperature measured in NH_3 (e.g. Lai et al. 2003). In the case of L1689B, for instance, the effective temperature of the Bonnor-Ebert fit shown in Fig. 1b is $T_{\text{eff}} \sim 50$ K, while the dust temperature observed with ISOPHOT is only $T_d \sim 11$ K (Ward-Thompson et al. 2002). Third, the physical process responsible for bounding the cores at some external pressure is unclear. These arguments suggest that prestellar cores cannot simply be described as isothermal hydrostatic structures and are either already contracting (e.g. Lee, Myers, Tafalla 2001) or experiencing extra support from static or turbulent magnetic fields (e.g. Curry & McKee 2000). It has also recently been pointed out that good Bonnor-Ebert fits can often be found for non-equilibrium, transient “cores” produced by turbulence (Ballesteros-Paredes et al. 2003).

As shown by Bacmann et al. (2000), one way to account for large density contrasts and high effective temperatures is to consider models of cores initially supported by a static magnetic field and evolving through ambipolar diffusion (e.g. Ciolek & Mouschovias 1994, Basu & Mouschovias 1995). A serious problem, however, with such models is that they need a strong field *in the low-density ambient cloud* (~ 30 – $100 \mu\text{G}$) to explain the observed sharp edges (see Bacmann et al. 2000). This seems inconsistent with existing Zeeman measurements (e.g. Crutcher 1999, Crutcher & Troland 2000). Furthermore, when the ambient field is strong (highly subcritical case), the model cores evolve on the ambipolar diffusion timescale $t_{AD} \sim 10 t_{ff}$ (Ciolek & Basu 2001; t_{ff} is the free-fall time), while observed cores have significantly shorter lifetimes $\sim 4 t_{ff}$ at $n_{\text{H}_2} \sim 10^4 \text{ cm}^{-3}$ (cf. Jessop & Ward-Thompson 2000). On the other hand, isolated starless cores appear to live $\gtrsim 3$ times *longer* than typical density fluctuations generated by turbulence (cf. Ballesteros-Paredes et al. 2003.) It is possible that more elaborate ambipolar diffusion models, incorporating the effects of a non-static, turbulent magnetic field in the outer parts of the cores and in the ambient cloud, would be more satisfactory and could also account for the filamentary shapes often seen on large

($\gtrsim 0.25$ pc) scales (cf. Jones & Basu 2002).

3. VELOCITY STRUCTURE OF CLASS 0 ENVELOPES

Another approach to constraining the collapse initial conditions consists in observing the detailed properties of very young accreting protostars (Class 0 objects – e.g. André, Ward-Thompson, Barsony 2000) that still retain detailed memory of their genesis. There have been very few quantitative studies of the velocity structure of the envelopes of Class 0 objects so far. Here, we describe two recent, contrasted examples.




Fig. 2. Position-velocity diagram along the major axis of the IRAM 04191 envelope (i.e., perpendicular to the flow) based on a $\text{C}^{34}\text{S}(2-1)$ map taken at the IRAM 30m telescope (Belloche et al. 2002). Contours: 0.2 to 0.8 by 0.2 K. The dots with error bars mark the observed velocity centroids. The solid curve shows the profile of a model with differential rotation [$V_{\text{rot}}(r > 3500 \text{ AU}) \propto r^{-1.5}$].

of the protostellar envelope does not occur in a rigid-body, but differential, fashion: the inner ~ 3500 AU-radius region rotates significantly faster than the outer parts of the envelope, as indicated by the characteristic “S” shape of the position-velocity diagrams obtained

The first object, IRAM 04191+1522 (IRAM 04191 for short), is relatively *isolated*. It was found at 1.3 mm by André, Motte, & Bacmann (1999) in the southern part of the Taurus cloud ($d = 140$ pc). IRAM 04191 is associated with a prominent, flattened protostellar envelope, seen in the (sub)millimeter dust continuum and in dense gas tracers such as N_2H^+ , C_3H_2 , H^{13}CO^+ , and DCO^+ (Belloche et al. 2002). All of the maps taken at the IRAM 30m telescope in small optical depth lines show a clear rotational velocity gradient across the envelope of $\sim 9 \text{ km s}^{-1} \text{ pc}^{-1}$ (after deprojection), perpendicular to the outflow axis. This gradient is one order of magnitude larger than those typically observed in starless cores (e.g. Goodman et al. 1993). Furthermore, the rotation




Fig. 3. Rotational velocity (a) and infall velocity (b) inferred in the IRAM 04191 envelope based on radiative transfer modeling of multi-transition CS and C^{34}S observations with the IRAM 30m telescope (Belloche et al. 2002). The shaded areas show the estimated domains where the models match the observations.

by Belloche et al. (see, e.g., Fig. 2). Fast, differential rotation is expected in protostellar envelopes because of conservation of angular momentum during dynamical collapse. In the present case, however, the dramatic drop in rotational velocity observed at $r \geq 3500$ AU (Fig. 3a), combined with the flat infall velocity profile (see below), points to *losses of angular momentum* in the outer envelope (see discussion in § 4 below).

Direct evidence for infall motions over a large portion of the envelope of IRAM04191 is observed in optically thick lines such as CS(2–1), CS(3–2), $\text{H}_2\text{CO}(2_{12} - 1_{11})$, and $\text{H}_2\text{CO}(3_{12} - 2_{11})$. These lines are double-peaked and skewed to the blue up to $\gtrsim 40''$ from source center, which is indicative of infall motions up to a radius $R_{\text{inf}} \gtrsim 5000$ AU (cf. Evans 1999, 2003). Radiative transfer modeling confirms this view, suggesting a flat, subsonic infall velocity profile ($V_{\text{inf}} \sim 0.1 \text{ km s}^{-1}$) for $3000 \lesssim r \lesssim 11000$ AU and larger infall velocities scaling as $V_{\text{inf}} \propto r^{-0.5}$ for $r \lesssim 3000$ AU (see Fig. 3b and Belloche et al. 2002 for details). The mass infall rate is estimated to be $\dot{M}_{\text{inf}} \sim 2 - 3 \times a_s^3/G \sim 3 \times 10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}$ (where $a_s \sim 0.15 - 0.2 \text{ km s}^{-1}$ is the sound speed at $T \sim 6 - 10 \text{ K}$), roughly independent of radius.

Another Class 0 object whose kinematics has been quantified in detail is IRAS 4A in the NGC 1333 *protocluster* (Di Francesco et al. 2001).

Using the IRAM Plateau de Bure interferometer, Di Francesco et al. observed inverse P-Cygni profiles in $\text{H}_2\text{CO}(3_{12} - 2_{11})$ toward IRAS 4A, from which they derived a very large mass infall rate of $\sim 1.1 \times 10^{-4} \text{ M}_{\odot} \text{ yr}^{-1}$ at $r \sim 2000 \text{ AU}$. Even if a warmer initial gas temperature ($\sim 20 \text{ K}$) than in IRAM 04191 and some initial level of turbulence are accounted for (see Di Francesco et al. 2001), this value of $\dot{M}_{\text{inf}}$ corresponds to more than ~ 15 times the canonical a_{eff}^3/G value (where $a_{\text{eff}} \lesssim 0.3 \text{ km s}^{-1}$ is the effective sound speed). This very high infall rate results both from a very dense envelope (a factor ~ 12 denser than a SIS at 10 K – see Motte & André 2001) and a large, supersonic infall velocity ($\sim 0.68 \text{ km s}^{-1}$ at $\sim 2000 \text{ AU}$ – Di Francesco et al. 2001). Evidence for fast rotation in the IRAS4A envelope, producing a velocity gradient as high as $\sim 40 \text{ km s}^{-1} \text{ pc}^{-1}$, was also reported by Di Francesco et al. (2001).

4. CONCLUSION: COMPARISON WITH COLLAPSE MODELS

In the case of *isolated dense cores* such as those of Taurus, the SIS model of Shu (1977) approximately describes some global features of the collapse (e.g. the mass infall rate within a factor ~ 2) but fails to reproduce the observations in detail. The extended infall velocity profiles observed in prestellar cores (Lee et al. 2001) and in the very young Class 0 object IRAM 04191 (Belloche et al. 2002 – see § 3) are indeed inconsistent with a pure inside-out collapse picture. The *shape* of the density profiles observed in prestellar cores are well fitted by purely thermal Bonnor-Ebert sphere models, but the *absolute* values of the densities are suggestive of some additional magnetic support (§ 2.1). The observed infall velocities are also marginally consistent with isothermal collapse models starting from Bonnor-Ebert spheres (e.g. Foster & Chevalier 1993, Hennebelle et al. 2003), as such models tend to produce somewhat faster velocities. This suggests that the collapse of ‘isolated’ cores is essentially *spontaneous* and somehow moderated by magnetic effects in (mildly) magnetized, non-isothermal versions of Bonnor-Ebert cloudlets. Indeed, the contrast seen in Fig. 3 between the steeply declining rotation velocity profile and the flat infall velocity profile of the IRAM 04191 envelope beyond $\sim 3500 \text{ AU}$ is very difficult to account for in the context of non-magnetic collapse models. In the presence of magnetic fields, on the other hand, the outer envelope can be coupled to, and spun down by, the (large moment of inertia of the) ambient cloud (e.g. Basu & Mouschovias 1995).

Based on a qualitative comparison with the ambipolar diffusion models of Basu & Mouschovias, Belloche et al. (2002) propose that the rapidly ro-

tating inner envelope of IRAM 04191 corresponds to a magnetically (slightly) supercritical core decoupling from an environment still supported by magnetic fields and strongly affected by magnetic braking. A magnetic field $\sim 60 \mu\text{G}$ is required at 3500 AU where $n_{\text{H}_2} \sim 1 - 2 \times 10^5 \text{ cm}^{-3}$, which is comparable to the field strengths recently estimated at such densities by Crutcher et al. (2003) in three prestellar cores. In this picture, the inner ~ 3500 AU radius envelope of IRAM 04191 would correspond to the effective mass reservoir ($\sim 0.5 M_\odot$) from which the central star is being built. Moreover, comparison of these results with the rotational characteristics of other objects in Taurus (Ohashi et al. 1997) indicates that IRAM 04191 behaves in a typical manner and is simply observed particularly soon after protostar formation. The IRAM 04191 example therefore suggests that *the masses of stars forming in clouds such as Taurus are largely determined by magnetic decoupling effects.*

In *protoclusters* such as NGC 1333, by contrast, the large overdensity factors measured in Class 0 envelopes compared to hydrostatic isothermal structures (Motte & André 2001), as well as the fast supersonic infall velocities and very large infall rates observed in some cases (§ 3), are inconsistent with self-initiated forms of collapse and require a *strong external influence*. This point is supported by the results of recent SPH simulations by Hennebelle et al. (2003, 2004). These simulations follow the evolution of a Bonnor-Ebert sphere whose collapse has been induced by an increase in external pressure P_{ext} . Large overdensity factors (compared to a SIS), together with supersonic infall velocities, and large infall rates ($\gtrsim 10 a_s^3/G$) are found near point mass formation when (and only when) the increase in P_{ext} is strong and very rapid (e.g. Fig. 4), resulting in a violent compression wave (cf. Whitworth & Summers 1985). Such




Fig. 4. Density profile (solid curve) obtained near point mass formation in SPH simulations of the collapse of an initially stable, rotating ($\beta = E_{\text{rot}}/E_{\text{grav}} = 2\%$) Bonnor-Ebert sphere ($T = 10$ K) induced by a very rapid increase in external pressure (with $P_{\text{ext}}/\dot{P}_{\text{ext}} = 0.03 \times$ the initial sound crossing time) (Hennebelle et al. 2004). Note the large overdensity factor compared to the $\rho \propto r^{-2}$ profile of a SIS at 10 K (dotted line).

a violent collapse in protoclusters may be conducive to the formation of both massive stars (through higher accretion rates) and multiple systems (when realistic, non-isotropic compressions are considered). Future high-resolution studies with the next generation of (sub)millimeter instruments (e.g., ALMA – Wootten 2001) will greatly help test this view and shed further light on the physics of collapse in cluster-forming regions.

REFERENCES

- ??lves, J.F., Lada, C.J., & Lada, E.A. 2001, *Nature*, 409, 159
- ??ndré, P., Motte, F., Bacmann, A. 1999, *ApJL*, 513, L57
- ??ndré, P., Ward-Thompson, D., Barsony, M. 2000, in *Protostars and Planets IV*, eds. V. Mannings, A.P. Boss, & S.S. Russell (Univ. of Arizona Press, Tucson), p. 59
- ??ndré, P., Ward-Thompson, D., Motte, F. 1996, *A&A*, 314, 625
- ??acmann, A., André, P., Puget, J.-L., Abergel, A., Bontemps, S., & Ward-Thompson, D. 2000, *A&A*, 361, 555
- ??allesteros-Paredes, J., Klessen, R.S., & Vázquez-Semadeni, E. 2003, *ApJ* 592, 188
- ??asu, S., & Mouschovias, T.Ch. 1995, *ApJ*, 453, 271
- ??elloche, A., Andr, P., Despois, D. & Blinder, S. 2002, *A&A*, 393, 927
- ??iolek, G.E., & Basu, S. 2001, in *From Darkness to Light*, Eds. T. Montmerle & P. Andr, ASP Conf. Ser., 243, p. 79
- ??iolek, G.E., Mouschovias, T.Ch. 1994, *ApJ*, 425, 142
- ??rutter, R.M. 1999, *ApJ*, 520, 706
- ??rutter, R.M., & Troland, T.H. 2000, *ApJ*, 537, L139
- ??rutter, R.M., Nutter, D.J., Ward-Thompson, D., & Kirk, J. et al. 2003, *ApJ* (astro-ph/0305604)
- ??urry, C.L., & McKee, C.F. 2000, *ApJ*, 528, 734
- ??i Francesco, J., Myers, P.C., Wilner, D.J. & Ohashi, N. 2001, *ApJ*, 562, 770
- ??vans, N.J. 1999, *ARA&A*, 37, 311
- ??vans, N.J. II 2003, in *Chemistry as a Diagnostic of Star Formation*, Eds. C.L. Curry & M. Fich, in press (astro-ph/0211526)
- ??oster, P.N., Chevalier, R.A. 1993, *ApJ* 416, 303 (FC93)

- Goodman, A.A., Benson, P.J., Fuller, G.A., Myers, P.C. 1993, *ApJ*, 406, 528
- Hennebelle, P., Whitworth, A.P., Gladwin, P.P., & Andr, P. 2003, *MNRAS*, 340, 870
- Hennebelle, P., Whitworth, A.P., Cha, S., Goodwin, S., 2004, *MNRAS*, in press (astro-ph/0311219)
- Jessop, N., & Ward-Thompson, D. 2000, *MNRAS*, 311, 63
- Johnstone, D., Wilson, C. D., Moriarty-Schieven, G., et al. 2000, *ApJ*, 545, 327
- Kanes, C.E., & Basu, S. 2002, *ApJ*, 569, 280
- Klessen, R.S., Heitsch, F., & Mac Low, M.-M. 2000, *ApJ*, 535, 887
- Kai, S.-P., Velusamy, T., Langer, W.D., & Kuiper, T.B.H. et al. 2003, *AJ*, 126, 311
- Kee, C.W., Myers, P.C., & Tafalla, M. 2001, *ApJS*, 136, 703
- Ki, Z.-Y., Shu, F.H. 1997, *ApJ*, 475, 237
- Kotte, F., André, P. 2001, *A&A*, 365, 440
- Kotte, F., André, P., Neri, R. 1998, *A&A*, 336, 150
- Kouschovias, T.C., & Ciolek, G.E. 1999, in *The Origin of Stars and Planetary Systems*, Eds. C.J. Lada & N.D. Kylafis, Kluwer, p.305
- Khashi, N., Hayashi, M., Ho, P.T.P., Momose, M., Tamura, M., Hirano, N., Sargent, A. 1997, *ApJ*, 488, 317
- Kadoan, P., & Nordlund, A. 2002, *ApJ*, 576, 870
- Kirley, Y., Evans II, N.J., Rawlings, J.M.C., Gregersen, E.M. 2000, *ApJS*, 131, 249
- Khu, F.H. 1977, *ApJ*, 214, 488
- Khu, F.H., Adams, F.C., Lizano, S. 1987, *ARA&A*, 25, 23
- Khu, F.H., Li, Z.-Y., Allen, A. 2003, preprint (astro-ph/0311426)
- Kiebenmorgen, R., & Krügel, E. 2000, *A&A*, 364, 625
- Kesti, L., Sargent, A.I 1998, *ApJL*, 508, L91
- Kard-Thompson, D., André, P., & Kirk, J.M. 2002, *MNRAS*, 329, 257
- Kard-Thompson, D., Motte, F., André, P. 1999, *MNRAS*, 305, 143
- Kard-Thompson, D., Scott, P.F., Hills, R.E., André, P. 1994, *MNRAS*, 268, 276
- Khitworth, A., Summers, D. 1985, *MNRAS*, 214, 1
- Kooten, A. 2001, *Science with the Atacama Large Millimeter Array*, ASP Conf. Ser., Vol. 235